

## PERIODONTAL DISEASE AND SYSTEMIC HEALTH

Lecturer Daniela Miricescu<sup>1</sup>, MD, PhD, Assoc. Prof. Alexandra Totan<sup>1</sup>, MD, PhD,  
Assist. Prof. Iulia-Ioana Stanescu<sup>1</sup>, MD, PhD, Assist. Prof. Radu Radulescu<sup>1</sup>, MD, PhD,  
Assist. Prof. Constantin Stefani<sup>2,3</sup>, MD, PhD, Assist. Prof. Ana Maria Alexandra Stanescu<sup>2</sup>, MD, PhD,  
Prof. Maria Greabu<sup>1</sup>, MD, PhD

<sup>1</sup> Department of Biochemistry, Faculty of Dental Medicine,

“Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

<sup>2</sup> Department of Family Medicine, “Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

<sup>3</sup> “Carol Davila” University Central Emergency Military Hospital, Bucharest, Romania

### ABSTRACT

Periodontitis is a chronic inflammatory disease of the oral cavity that usually affects the adult population. In the last two decades, many specialized studies have reported that there is a link between periodontitis, cardiovascular diseases and diabetes mellitus. Main factors that lead to periodontal disease installation are several bacterial species that induce both local and systemic inflammation, negatively contributing to the progression of cardiovascular diseases. One of the many complications of diabetes mellitus is periodontitis, there is a two-way relationship between these two diseases. The purpose of this review is to analyze the recent data provided by the literature regarding the relationship between oral periodontal pathogens and systemic health.

**Keywords:** bacterial pathogens, inflammation, diabetes mellitus, cardiovascular diseases

### PERIODONTAL DISEASE PATHOGENESIS

Periodontal diseases are divided into two groups: reversible and irreversible. Gingivitis is a reversible inflammatory condition due to dental plaque accumulation, affecting approximately 50% of the adult population. It is characterized by an initial increase in blood flow, increased vascular permeability and an influx of cells (polymorphonucleated leukocytes (PMNs) and macrophages) from the peripheral blood to the periodontal connective tissue (1-3).

Periodontal disease is the most common chronic inflammation of the oral cavity in the adult population worldwide. 35% of the adult population worldwide has periodontitis, 11% have a moderate or severe form, this oral disease being considered a pandemic health problem that causes disabilities, speech impairment, low self-esteem and reduced quality of life (4,5).

This chronic inflammatory disease affects the supporting tissues of the tooth, leading to the pro-

gressive destruction of connective tissue and alveolar bone (5,6). After initiation, the disease progresses with the loss of collagen fibers and their disintegration from the cement surface, apical migration of epithelial cells, formation of periodontal pockets and resorption of alveolar bone. If left untreated, the disease continues with the progressive destruction of the bone, leading to tooth mobility and finally tooth loss (5-7).

The role of microorganisms in the etiology of periodontitis has been very well studied, so between 500-700 bacteria species are able to colonize oral cavity. 10% of these bacteria plays important roles in the initiation of periodontal disease. Periodontitis is initiated by colonization of the gums by pathogenic bacteria such as *Porphyromonas gingivalis*, *Actinobacillus actinomycetemcomitans* and *Bacteroides forsythicus*. *P. gingivalis*, *Tannerella forsythensis* and *Troponema denticola*, are directly associated with chronic periodontitis. *A. actinomycetemcomitans* has been observed in the early

Corresponding author:

Assoc. Prof. Alexandra Totan, “Carol Davila” University of Medicine and Pharmacy, Bucharest, Romania

E-mail: alexandratotan99@gmail.com

stages of periodontal disease and in aggressive periodontitis (8,9).

These bacterial species have the ability to activate local defense mechanisms, destroy the epithelium and other structures of the gingiva, while inactivating their repair systems. Activated neutrophils present at the site of inflammation are immobilized and produce reactive oxygen species (SRO) that are capable of destroying the pathogens. ROS overproduction can damage the surrounding host tissue (8-10).

There are a number of systemic factors that can contribute to the progression of periodontal diseases such as heredity, nervous system disorders (psychicstress), endocrine imbalances (diabetes), immune deficiencies, cardiovascular, hematological, hepatic, renal diseases and nutrition deficient (2,11).

Are three important biological stages in the progression of periodontal disease: inflammation, connective tissue degradation and the last one bone turnover. Inflammation in periodontal disease is response to bacterial plaque accumulation, permanent presence of bacterial multispecies, leads to chronic inflammation and abundance of oral inflammatory biomarkers (6,8,9). Many studies have detected various inflammation biomarkers to be increased both saliva or gingival crevicular fluid (GCF) in patients with periodontitis. Interleukin-1 (IL-1), IL-6, C-reactive protein (CRP) tumor necrosis factor-alpha (TNF- $\alpha$ ), prostaglandin E2 are specific biomarkers to the inflammatory process which have been detected both in total unstimulated saliva and also CGF, where they present increased levels (6,9, 12-17). Reactive oxygen species (ROS) can be generated by inflammatory reactions, especially chronic ones. Inflammatory cells such as activated neutrophils and macrophages release different types of ROS such as  $O_2^{\cdot-}$ ,  $O_2^{\cdot-}$ ,  $H_2O_2$ ,  $HO^\circ$  and  $HOCl$  (18). The production of ROS by the PMN under inflammatory conditions ensures a protective role for the host. Overproduction of ROS by PMNs, leads to oxidative damage of gingival tissue, periodontal ligament and alveolar bone (19-21).

8-hydroxydeoxyguanine (8-HOdG-biomarker of DNA damage), malondialdehyde (MDA -marker of lipidperoxidation) were detected to be increased in saliva and serum of patients with chronic periodontitis. At the same time, statistically lower levels of uric acid, catalase and glutathione peroxidase were detected in saliva and serum of patients with chronic *versus* healthy volunteers (20-23).

Matrixmetalloproteases (MMPs) are local proteinases responsible for both degradation and remodeling. During periodontal disease evolution, collagen fibers are destroyed by interstitial collagenases, derived from the host cells. A collagenase capable of degrading the structure of the triple helix of type I, II, III collagen in the alveolar bone matrix is collagenase 2 or MMP-8. MMP-8 is released in increased levels when the PMNs are mobilized to inflammation site (23,24). Increased salivary or GCF levels of MMP-8, myeloperoxidase (MPO) and tissue inhibitor of matrix metalloproteinases (TIMP)-1 were detected in patients with periodontal disease. Osteocalcin is a specific marker of bone, his release may induce a new osteoid synthesis or resorption of the alveolar bone. Increased salivary statistically levels of osteocalcin was observed in patients with periodontal disease (23,25). Type I collagen C-terminal telopeptide (CTX) is one of the most studied members of collagen degradation molecules, which present an increased salivary level in patients with chronic periodontitis (23). An organism's response to periodontal infection involves the release of enzymes from stromal, epithelial, inflammatory or bacterial cells. Detection of these enzymes in saliva and GCF may help to clarify the pathogenesis of periodontal disease. Important roles in tissue degradation are played by enzyme ssuch as: elastases, gelatinases and proteinases. Aspartate aminotransferase (AST), lactate dehydrogenase (LDH), alkaline phosphatase (ALP), acid phosphatase (ACP), glutamyl transferase (GGT) and creatine kinase (CK) are intracellular enzymes that are released in a large amount from the injured cells of periodontal tissues in FCG and saliva as well as in surrounding fluid. These enzymes usually present increased levels in patients diagnosed with periodontal disease (26,27).

## PERIODONTAL DISEASE AND DIABETES MELLITUS

Between diabetes and periodontal disease has been observed a bidirectional relationship, in which diabetes increases the risk of developing periodontal disease and periodontal inflammation negatively affects glycemic control. Diabetes mellitus is a chronic metabolic disorder that affects the population worldwide, which can be characterized either by the lack of insulin production by the pancreatic beta cells (type 1 diabetes), or by insulin resistance (type 2 diabetes), conditions that will leads to hyperglycemia (28,29).

Diabetes mellitus has a number of well-known complications such as retinopathy, nephropathy, poor wound healing, neuropathy, macro and microvascular impairment and periodontal disease (29,30). An observational study conducted by De Miguel-Infants and coworkers in adults aged 40 or older, observed that diabetic patients have an increased risk of developing periodontal disease (31).

Chronic inflammation that is specific during the periodontal progression causes an exacerbated inflammatory response correlated with low metabolic control of serum glucose and increased insulin requirements (32). In patients with type 2 diabetes mellitus, Aikten JP and coworkers detected a positive correlation between salivary levels of alpha-2-macroglobulin and HbA1C, this aspect reflecting that the glycemic control of patients can also be performed at the salivary level (33). The salivary concentration of melatonin has been detected to be decreased in patients with type 2 diabetes mellitus and in patients with periodontal disease, which may be a key biomarker in the diagnosis and treatment of the two diseases (34). Patients with gingivitis or periodontitis, diabetic patients, present increased levels of glucose,  $\alpha$ -hydroxybutyrate and biomarkers of oxidative stress (35). In a study conducted by Liu and coworkers on diabetic mice it was observed that periodontal disease aggravated pancreatic  $\beta$ -cell failure and induces insulin resistance (36). Individuals with acute bacterial or viral infection are unfortunately characterized by severe and long-lasting insulin resistance. This aspect being confirmed by Fernandez-Real JM and coworkers in a study of 124 infected middle-aged men with enteroviruses and *C. pneumonia* (37).

Sugiyama S et al. observed in a diabetic rats study infected with *P. gingivalis* a decrease in gingival vascular function and increase in insulin resistance (38). Diabetic patients have a 3-fold higher risk of developing periodontitis compared to healthy individuals (39).

An incidence of periodontal disease of 58% was observed in patients with type 1 diabetes, compared with non-diabetic patients, who had a 15% incidence (40). Diabetes mellitus causes morphological changes in the salivary glands and in saliva composition, so Malicka B and coworkers observed in diabetic patients that myeloperoxidase and salivary Ig A were correlated with poor periodontal status (41). Increased serum levels of inflammation mediators such as CRP, TNF- $\alpha$ , and IL-6 were detected in both patients with periodontal disease and diabetic patients (42-44). OS is a major link be-

tween the 2 diseases, plasma and salivary levels of OS have been detected to be increased in both conditions that further can activate pro-inflammatory systemic pathways (45).

## PERIODONTITIS AND CARDIOVASCULAR DISEASES

Cardiovascular diseases is a major cause of death worldwide including in Romania. Diabetes, smoking and inflammation are risk factors for both myocardial infarction and periodontal disease (46-50). Have been observed a possible association between cardiovascular diseases and periodontitis, but the role of oral infection at the cardiovascular level is not fully elucidated. The results of cohort studies observed that people with spinal cord disease have a 2.22-fold higher risk of developing cardiovascular diseases (51). Pussinen PJ and coworkers detected at 1163 men patients with coronary heart disease in serum antibodies against *P. gingivalis*, and *A. actinomycetemcomitans* (52).

Figuro E and coworkers detected in 42 atheromatous plaques, bacteria specific to periodontal pathology such as *P. gingivalis*, *A. actinomycetemcomitans*, *T. forsythia*, but also other bacterial types such as *Eikenella corrodens*, *Fusobacterium nucleatum* and *Campylobacter rectus* (53). The results of this study confirm that oral periodontal pathogens may migrate to other body sites. Studies in mice receiving a hyperlipidic diet to develop atheromatous plaque and infected with *P. gingivalis* and *T. denticola* have shown that the presence of these bacteria is associated with alveolar bone loss and aortic atherosclerosis (54).

*P. gingivalis* and *T. denticola* can induce a systemic immune response, their bacterial genomic DNA being detected in the oral epithelia and aorta. *P. gingivalis* being the only oral bacterial species that induces the expression of virulent factors associated with platelet aggregation (55). Salivary amylase may be an independent diagnostic factor for acute myocardial infarction in patients suffering from precordial pain less than 4 hours.

The use of nano-biochips as a screening method based on salivary proteins such as CRP, myoglobin and myeloperoxidase may be beneficial for patients suffering from acute myocardial infarction (56,57).

Several studies have suggested a possible connection between periodontal disease and endothelial dysfunction, caused by reduced bioavailability of nitric oxide, that reduces platelet aggregation, inhibits leukocyte attachment to endothelial cells,

and prevents the expression of adhesion molecules (58,59). Studies on rats with periodontitis confirmed the reduction of nitric oxide level and the correlation with endothelial dysfunction (59-61).

Endotoxins and antigens secreted by periodontal pathogens contribute to the pathogenesis of endothelial dysfunction, by stimulating the attachment of leukocytes to the surface of endothelial cells, may induce the expression of adhesion molecules such as endothelial monocytes chemoattractant protein -1 (MCP-1), may induce a significant rise and release of endothelin-1 (62,63). Has been observed that periodontal treatment supplemented with antibiotics improve endothelial dysfunction (64). Stroke presents a number of well-known risk factors such as hypertension, dyslipidemia, diabetes, smoking and age. Numerous epidemiological studies have also included periodontal disease as a major risk factor (65,66). Increased serum levels of CRP and antibodies against *P. gingivalis* have been associated with acute ischemic stroke (66,67). Per-

iodontal pathogens can be found in heart valve tissue, atrial and ventricular tissues, can colonize atherosclerotic plaque (68,69).

**In conclusion**, periodontal pathogens or their endotoxins induce an inflammatory response both locally and systemically. *P. gingivalis* is the most aggressive periodontal pathogen at cardiac level being able to induce platelet aggregation and the expression of numerous adhesion molecules contributing in this way to the progression of cardiovascular diseases. Children and adults who have been diagnosed with type 1 and type 2 diabetes have a rather high risk of developing periodontal disease. Periodontal disease should not be ignored and proper treatment will greatly improve the patients health both locally and systemic.

### Acknowledgement

All authors equally contributed to the present paper.

### REFERENCES

- Ridgeway EE. Periodontal disease: diagnosis and management. *J Am Acad Nurse Pract* 2000; 12:79-83.
- Miyata Y, Obata Y, Mochizuk Y et al. Periodontal Disease in Patients Receiving Dialysis. *Int J Mol Sci* 2019; 20: 3805:1-21.
- Maderal AD, Salisbury L, Jorizzo JL. Desquamative gingivitis: Diagnosis and treatment. *J Am Acad Dermatol* 2018; 78(5):851-861.
- Richards D. Oral diseases affect some 3.9 billion people. *Evid Based Dent* 2013; 14:35.
- Peres MA, Macpherson LMD, Weyant RJ. Oral diseases: a global public health challenge. *Lancet* 2019; 20: 394(10194):249-260.
- Rai B, Kaur JM, Kharb S et al. Peripheral blood and C-reactive protein levels (CRP) in chronic periodontitis. *African J Biochem Res* 2009; 3(4): 150-153.
- Miller CS, Foley JD, Bailey AL et al. Current developments in salivary diagnostics. *Biomarkers Med* 2010; 4(1): 1-19.
- Jia L, Han N, Du J et al. Pathogenesis of Important Virulence Factors of *Porphyromonas gingivalis* via Toll-Like Receptors. *Front Cell Infect Microbiol* 2019; 18:9:262.
- Kinney J, Ramseier C, Giannobile WV. Oral fluid-based biomarkers of alveolar bone loss in periodontitis. *Ann N Y Acad Sci* 2007; 1098:230-251.
- Dewhirst FE, Chen T, Izard J et al. The human oral microbiome. *J Bacteriol* 2010; 192: 5002-5017.
- Falcao A, Bullón P. A review of the influence of periodontal treatment in systemic diseases. *Periodontol 2000* 2019;79(1):117-128.
- Konopka T, Krol K, Kopec W et al. Total antioxidant status and 8-hydroxy-2'-deoxyguanosine levels in gingival and peripheral blood of periodontitis patients. *Arch Immunol Ther Exp* 2007; 55: 417-425.
- Machado-Moreira EA, Filho DW, Bittencourt de Oliveira T et al. Proinflammatory and oxidative stress markers in patients with periodontal disease. *Mediators Inflamm* 2007; 45794.
- Vidita-Gurban C, Suci O, Vlaia L et al. Inflammatory Markers from Crevicular Fluid in Periodontal Disease. *Rev. Chimie* 2016; 67(10):1952-1956.
- Miller CM, King CP, Chris-Langub Jr M et al. Salivary biomarkers of existing periodontal disease: A cross-sectional study. *J Am Dent Assoc* 2006;137: 322-329.
- Rai B, Kaur J, Anand SC et al. Salivary stress markers, stress and periodontitis: a pilot study. *J Periodontol* 2011; 82(2): 287-92.
- Braz-Silva PH, Bergamini ML, Mardegan AP et al. Inflammatory profile of chronic apical periodontitis: a literature review. *Acta Odontol Scand* 2019;77(3): 173-180.
- Valko M, Leibfritz D, Moncol J et al. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol* 2007; 39: 44-84.
- Da Silva JC, Muniz FWMG, Oballe HJR et al. The effect of periodontal therapy on oxidative stress biomarkers: A systematic review. *J Clin Periodontol* 2018 ;45(10):1222-1237.
- Scully DV, Langlez-Evans SC. Salivary antioxidants and periodontal disease status. *Proc Nutr Soc* 2002; 61:137-143.
- Tóthová L, Celec P. Oxidative Stress and Antioxidants in the Diagnosis and Therapy of Periodontitis. *Front Physiol* 2017;8:1055.
- Canakci CF, Cicek Y, Yildirim A et al. Increased levels of 8-hydroxydeoxyguanosine and malondialdehyde and its relationship with antioxidant enzymes in saliva of periodontitis patients. *Eur J Dent* 2009; 3:100-106.
- Miricescu D, Totan A, Calenic B et al. Salivary biomarkers: relationship between oxidative stress and alveolar bone loss in chronic periodontitis. *Acta Odontol Scand* 2014; 72(1):42-47.
- Romero-Castro NS, Vázquez-Villamar M, Muñoz-Valle JF et al. Relationship between TNF- $\alpha$ , MMP-8, and MMP-9 levels in gingival crevicular fluid and the subgingival microbiota in periodontal disease. *Odontology* 2019; doi: 10.1007.
- Nascimento GG, Baelum V, Sorsa T et al. Salivary levels of MPO, MMP-8 and TIMP-1 are associated with gingival inflammation response patterns during experimental gingivitis. *Cytokine* 2019;115:135-141.
- Todorov T, Dozic I, Barrero MV et al. Salivary enzymes and periodontal disease. *Med Oral Patol Oral Circ Bucal* 2006;11: E115-119.
- Greabu M, Miricescu D, Totan A et al. Salivary and serum enzymes as diagnostic biomarkers in patients with periodontal disease. *Stoma Edu J* 2014; 1(1): 22-26.

28. Salmerón D, Gómez García F, Pons-Fuster et al. Screening for prediabetes and risk of periodontal disease. *Diabetes Metab Syndr* 2019;13(2):1661-1666.
29. Stanko P, Izakovicova H L. Bidirectional association between diabetes mellitus and inflammatory periodontal disease. A review. *Biomed Pap Med Fac Univ Palacky* 2014; 158: 35-38.
30. Laiginhas R, Madeira C, Lopes Met et al. Risk factors for prevalent diabetic retinopathy and proliferative diabetic retinopathy in type 1 diabetes. *Endocrine* 2019;12. doi: 10.1007.
31. De Miguel-Infante A, Martinez-Huedo MA, Mora-Zamorano E et al. Periodontal disease in adults with diabetes, prevalence and risk factors. Results of an observational study. *Int J Clin Pract* 2018;e13294.
32. Nishimura F, Iwamoto T, Mineshiba J et al. Periodontal disease and diabetes mellitus: the role of tumor necrosis factor- $\alpha$  in a 2-way relationship. *J Periodontol* 2003; 74:97-102.
33. Aitken JP, Ortiz C, Morales-Bozo I et al.  $\alpha$ -2-macroglobulin in saliva is associated with glycemic control in patients with type 2 diabetes mellitus. *Dis Markers* 2015; 2015: 128653.
34. Abdolsamadi H, Goodarzi MT, Motemayel FA et al. Reduction of melatonin level in patients with type II diabetes and periodontal diseases. *J Dent Res Dent Clin Dent Prospects* 2014; 8(3): 160-165.
35. Satish BN, Srikala P, Maharudrappa B et al. Saliva: a tool in assessing glucose levels in diabetes mellitus. *J Int Oral Health* 2014; 6(2): 114-117.
36. Liu Y, Zhang QJ. Periodontitis aggravated pancreatic  $\beta$ -cell dysfunction in diabetic mice through interleukin-12 regulation on Klotho. *Diabetes Investig* 2016;7: 303-311.
37. Fernandez-Real JM, Lopez-Bermejo A, Vendrell J et al. Burden of infection and insulin resistance in healthy middle-age men. *Diabetes Care* 2006;29:1058-1064.
38. Sugiyama S, Takahashi SS, Tokutomi FA et al. Gingival vascular functions are altered in type 2 diabetes mellitus model and/or periodontitis model. *J Clin Biochem Nutr* 2012; 51:108-113.
39. Mealey BL, Ocampo GL. Diabetes mellitus and periodontal disease. *J Periodontol* 2006; 77: 1289-1303.
40. Poplawska-Kita A, Siewko K, Szpak P et al. Association between type 1 diabetes and periodontal health. *Adv Med Sci* 2014; 59:126-131.
41. Malicka B, Kaczmarek U, Skoskiewicz-Malinowska K. Selected antibacterial factors in the saliva of diabetic patients. *Arch Oral Biol* 2015;60:425-431.
42. Engebretson S, Chertog R, Nichols A et al. Plasma levels of tumor necrosis factor- $\alpha$  in patients with chronic periodontitis and type 2 diabetes. *J Clin Periodontol* 2007; 34: 18-24.
43. Noack B, Genco RJ, Trevisan M et al. Periodontal infections contribute to elevated systemic C-reactive protein level. *J Periodontol* 2001; 72, 1221-1227.
44. Chen L, Wei B, Li J et al. Association of periodontal parameters with metabolic level and systemic inflammatory markers in patients with type 2 diabetes. *J Periodontol* 2010; 81: 364-371.
45. Patil VS, Patil VP, Gokhale N et al. Chronic Periodontitis in Type 2 Diabetes Mellitus: Oxidative Stress as a Common Factor in Periodontal Tissue Injury. *J Clin Diagn Res* 2016; 10: 12-16.
46. Diaconu CC, Stănescu AMA, Stoian AP et al. Hyperkalemia and cardiovascular diseases: new molecules for the treatment. *Rev Chim* 2018;69(6): 1367-1370.
47. Kjellström B, Gustafsson A, Nordendal E et al. Symptoms of depression and their relation to myocardial infarction and periodontitis. *Eur J Cardiovasc Nurs* 2017; 16: 468-474.
48. Stănescu AMA, Grajdeanu IV, Șerban B et al. Distribution of Cardiovascular Risk Factors in Patients with Psoriasis. *Rev Chim* 2019;70(5):1643-1648.
49. Shrihari TG. Potential correlation between periodontitis and coronary heart disease – an overview. *Gen Dent* 2012; 60: 20-24.
50. Bejan GC, Stănescu AMA, Ghilencea NL et al. Analiza actuală a aspectelor terapeutice moderne în tratamentul HTA și al comorbidităților asociate. *RMJ* 2018;2:102-107.
51. Bahekar AA, Singh S, Saha S et al. The prevalence and incidence of coronary heart disease is significantly increased in periodontitis: a meta-analysis. *Am Heart J* 2007; 154:830-837.
52. Pussinen PJ, Jousilahti P, Alfthan G et al. Antibodies to periodontal pathogens are associated with coronary heart disease. *Atheroscler Tromb Vasc Biol* 2003;23:1250-1254.
53. Figuro E, Sanchez-Beltran M, Cuesta-Frechoso S et al. Detection of periodontal bacteria in atheromatous plaque by nested polymerase chain reaction. *J Periodontol* 2011; 82:1469-1477.
54. Chukkappalli SS, Rivera MF, Velsko IM et al. Invasion of oral and aortic tissues by oral spirochete *Treponema denticola* in apo E(-/-) mice causally links periodontal disease and atherosclerosis. *Infect Immun* 2014; 82:1959-1967.
55. Sharma A, Novak EK, Sojar HT et al. Porphyromonas gingivalis platelet aggregation activity: outer membrane vesicles are potent activators of murine platelets. *Oral Microbiol Immunol* 2000; 15: 393-396.
56. Shen YS, Chen WL, Chang HY et al. Diagnostic performance of initial salivary alpha-amylase activity for acute myocardial infarction in patients with acute chest pain. *J Emerg Med* 2012;43:553-560.
57. Floriano PN, Christodoulides N, Miller CS et al. Use of saliva-based nano-biochip tests for acute myocardial infarction at the point of care: a feasibility study. *Clin Chem* 2009;55:1530-1538.
58. Garg UC, Hassid A. Nitric oxide-generating vasodilators and 8-bromo-cyclic guanosine monophosphate inhibit mitogenesis and proliferation of cultured rat vascular smooth muscle cells. *J Clin Invest* 1989; 83: 1774.
59. Punj A, Shenoy SB, Subramanyam K. Comparison of Endothelial Function in Healthy Patients and Patients With Chronic Periodontitis and Myocardial Infarction. *Periodontol* 2017;88(12):1234-1243.
60. Brito LC, DalBó S, Striechen TM et al. Experimental periodontitis promotes transient vascular inflammation and endothelial dysfunction. *Arch Oral Biol* 2013; 58: 1187-1198.
61. Campi P, Herrera BS, de Jesus FN et al. Endothelial dysfunction in rats with ligature-induced periodontitis: Participation of nitric oxide and cyclooxygenase-2-derived products. *Arch Oral Biol* 2016; 63: 66-74.
62. Hashizume T, Kurita-Ochiai T, Yamamoto M. Porphyromonas gingivalis stimulates monocyte adhesion to human umbilical vein endothelial cells. *FEMS Immunol Med Microbiol* 2011; 62: 57-65.
63. Vanhoutte PM, Shimokawa H, Feletou M et al. Endothelial dysfunction and vascular disease: a 30th anniversary update. *Acta Physiol* 2017; 219: 22-96.
64. Tonetti MS, D'Aiuto F, Nibali L et al. Treatment of periodontitis and endothelial function. *N Engl J Med* 2007; 356: 911-920.
65. Benjamin EJ, Muntner P, Alonso A et al. American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2019 Update: A Report From the American Heart Association. *Circulation* 2019; 139: e56-e66.
66. Leira Y, Seoane J, Blanco M et al. Association between periodontitis and ischemic stroke: A systematic review and meta-analysis. *Eur J Epidemiol* 2017; 32: 43-53.
67. Hosomi N, Aoki S, Matsuo K et al. Association of serum anti-periodontal pathogen antibody with ischemic stroke. *Cerebrovasc Dis* 2012; 34: 385-392.
68. Kozarov EV, Dorn BR, Shelburne CE et al. Human atherosclerotic plaque contains viable invasive *Actinobacillus actinomycetemcomitans* and *Porphyromonas gingivalis*. *Arterioscler Thromb Vasc Biol* 2005; 25: e17-e18.
69. Snow DE, Everett J, Mayer G et al. The presence of biofilm structures in atherosclerotic plaques of arteries from legs amputated as a complication of diabetic foot ulcers. *J Wound Care* 2016; 25: S16-S22