

Exploring psychosocial challenges  
among patients with sickle cell disease:  
A cross-sectional study in Basra city,  
Iraq

*By Zainab Barakat*

# Exploring psychosocial challenges among patients with sickle cell disease: A cross-sectional study in Basra city, Iraq

Qutaiba M. Dawood<sup>1</sup>, Zainab Barakat<sup>2</sup>, Amenah S. Abdulkareem<sup>1</sup>, Mustafa A. Jabr<sup>3</sup>, Murtaza H. Abd-Alsada<sup>3</sup>, Sara A. Sahb<sup>3</sup>, Maryam A. Zahru<sup>3</sup>, Zahra S. Maran<sup>3</sup>

1 Clinical Hematology, Department of Internal Medicine, Al Zahraa College of Medicine, University of Basrah, Basrah, Iraq.

2 Department of Medicine, College of Medicine, University of Basrah, Basrah, Iraq.

3 Al Zahraa College of Medicine, University of Basrah, Basrah, Iraq.

*Corresponding author:*

Zainab Barakat

*E-mail:* za4389452@gmail.com

## ABSTRACT

**Background:** Sickle cell disease (SCD) is a group of inherited hemoglobin disorders marked by alterations affecting the  $\beta$ -globin chain of hemoglobin. These could lead to persistent hemolytic anemia, severe acute and chronic pain. The study aimed to comprehend the impact of SCD on various facets of patients' lives, including education, employment, relationships, and mental health.

**Methods:** This descriptive cross-sectional study examines the psychosocial challenges of sickle cell disease (SCD) among patients in Basra city, Iraq. Conducted over three months at the Basra Hematology Center in Al Sayyab Teaching Hospital, the study enrolled 120 patients aged 15 and above diagnosed with SCD or sickle thalassemia. Utilizing structured interviews and standardized questionnaires.

**Results:** The prevalence of psychosocial challenges, with a significant proportion reporting difficulties in education, career choices, social relationships, and mental well-being. Pain intensity assessments using the Visual Analog Scale underscored the severity of pain experienced by participants. Additionally, socio-demographic factors such as age showed associations with pain scores.

Conclusion: This study highlights the significant psychosocial challenges faced by SCD patients, affecting various aspects of their lives including education, employment, relationships, and mental health. While certain socio-demographic factors showed associations with pain scores.

**Keywords:** sickle cell disease, psychosocial challenges, chronic pain, vasoocclusive crises,  $\beta$ -globin chain

8

## INTRODUCTION

Sickle cell disease (SCD) is a group of inherited hemoglobin disorders marked by alterations affecting the  $\beta$ -globin chain of hemoglobin, leading to persistent hemolytic anemia, severe acute and chronic pain, and ongoing organ damage [1,2]. Pain is a distinctive characteristic of SCD, significantly impacting patients' well-being and quality of life [3]. It manifests in various forms including acute intermittent pain, chronic daily pain, and acute-on-chronic pain, persisting from early infancy throughout patient's life [4]. As individuals age, chronic pain becomes increasingly prevalent, affecting 30% to 40% of SCD patients [5]. Vasoocclusive crises triggered by SCD often lead to intense acute pain, resulting from blood flow obstruction by sickled red blood cells in small vessels, reducing oxygen supply to affected organs. This pain can affect various anatomical regions, including bone structure, joints, and soft tissue, presenting symptoms such as dactylitis, acute joint necrosis, or acute abdomen [6]. Chronic pain in SCD refers to persistent pain experienced on most days for at least six months, and potentially originating from central sensitization or neuropathic mechanisms [7, 8]. The pain experienced in sickle cell disease often necessitates hospitalization, negatively impacts quality of life, potentially reduces survival rates, and leads to significant disability for many individuals [6]. Assessment of pain in SCD relies on patient self-reports, corroborated to identify underlying causes such as avascular necrosis or leg ulcers, with validated instruments crucial for clinical evaluation and decision-making [9].

In addition to the physical illness, sickle cell disease (SCD) patients also encounter a spectrum of psychosocial challenges, including social stigmatization, mental health disorders,

sleep disruptions, interpersonal dynamics, substance dependency, and workplace biases [10]. Depression and anxiety frequently affect SCD patients, resulting in high pain levels and clinical complications, suboptimal therapy adherence, and disruptions in various life aspects [10,11]. Another issue is social stigmatization that stems from diverse sources, including the use of opioids for pain management, racial identity (especially being of African descent), delayed growth or puberty, socioeconomic status, and the severity of the illness. Those with SCD who face stigma often contend with impaired sexual function, elevated stress levels, increased pain perception, inadequate coping mechanisms, medication non-adherence, and depression [12]. Moreover, sleep disturbances is also a common challenge. Notably, patients often experience a negative association between pain levels and sleep quality [13]. The most common sleep-disordered breathing is obstructive sleep apnea (OSA) which is attributed to upper airway obstruction resulting from compensatory adenotonsillar hyperplasia. This condition follows splenic infarction and reactive lymph node enlargement due to recurrent infections [14].

SCD also poses a great risk for developing depression and anxiety, as demonstrated in a recent study by Al Marzouki et al. [15] involving 119 patients aged 18 and older that concluded that 45.4% of participants exhibited depressive symptoms, with slightly higher rates among male patients. Depression and anxiety can also exacerbate pain and hinder coping strategies, with reported incidences ranging from 26% to 33% for depression and 6.5% to 36% for anxiety [9]. This study aimed to explore the psychosocial experiences and pain intensity among individuals aged 15 and above with SCD.

## **METHODS**

### **Study design**

This descriptive cross-sectional study was conducted in Basra city at the Basra Hematology Center in Al Sayyab Teaching Hospital, spanning three months from September 1st to November 31st, 2023. The study was conducted three days a week, with each session lasting two hours from 10:30 a.m. to 12:30 p.m.

### **Sampling, methods and sample size**

A convenient sample of 120 patients aged 15 and above, diagnosed with sickle cell disease (SCD) or sickle thalassemia, and attending the Hematology Center at Al-Sayyab Teaching Hospital in Basra city, were included in the study. The sample size was determined based on the available time period for the study. Patients included in the study either visited Basra Hematology Center independently or were referred from hematological clinics. The authors provided clear explanations regarding the purpose and nature of the study to all participants, emphasizing confidentiality, and obtained verbal consent from each subject before enrollment. Each selected patient underwent an interview in a private setting at the hospital using a specially designed structured questionnaire (see Appendix.2) tailored for the study's objectives.

### **Interviews**

The interviews lasted approximately 10-15minutes and aimed to comprehensive understanding of the psychosocial encounters of SCD patients and how pain affects their daily activities.

### **Questionnaire**

Patients who consented to participate completed standardized questionnaires in a private hospital setting. A medical student was present to assist with questionnaire administration and address any questions. A researcher-designed questionnaire was utilized. It comprised two sections: Section A focused on gathering respondents' demographic information, while Section B consisted of 20 items aimed at measuring the psychosocial challenges faced by SCD patients. The questionnaire's validity was ensured through expert validation in family medicine, while its reliability was established via the test-retest method, yielding a coefficient of 0.84.

### **Pain intensity**

**1** Pain was assessed using the Visual Analog Scale (VAS), a method used to gauge pain severity by asking patients to rate their pain level **11** on a scale from 0 to 10. Here, 0 denotes no pain, while 10 represents the most severe pain imaginable. Based on their scores, patients were categorized as follows:

- 0 = No pain.
- 1-2 = Mild, bothersome pain.
- 3-4 = Persistent, uncomfortable, bothersome pain.
- 5-6 = Distressing, distressing, distressing pain.
- 7-8 = Intense, dreadful, excruciating pain.
- 9-10 = Worst possible, unbearable, excruciating pain.

### **Data Collection Procedure**

Identification and Recruitment: Potential participants were identified by the authors at Basra Hematology Center. They were approached and informed about the study's objectives and procedures.

### **Data Management**

The research team securely stored and managed completed questionnaires and interview transcripts.

### **Informed Consent**

Every participant received detailed information about the study, including their rights as participants, the questionnaire, and the confidentiality protocols for their data. Informed consent was obtained from each participant.

### **Confidentiality**

Personal information and responses from participants were treated with utmost confidentiality and were securely stored and accessible only to authorized researchers.

### **Inclusion Criteria**

Patients diagnosed with SCD/sickle thalassemia, aged 15 and above, of both sexes, attending Basra Hematology Center.

### **Exclusion Criteria:**

1. Patients diagnosed with SCD under the age of 15.
2. Patients with sickle cell trait, pregnant individuals, and those diagnosed with hematological malignancies.
3. Post-operative SCD patients.
4. Patients who declined participation in the study.

**Statistical Analysis:** SPSS program (Statistical Packages for Social Sciences- version 19) <sup>4</sup> was used to carry out statistical analysis of the data. The results were tabulated. Chi-square test ( $\chi^2$ ) or

Fisher exact test, coupled with logistic regression analysis, were employed to discern associations among variables. Statistical significance was established with a p-value below 0.05.

## RESULTS


The socio-demographic features of the population are shown in Table.1. This study included 120 participants, with ages ranging from 15 to 65 years. The majority fell within the 15-24 age group, accounting for 55.8%. Those aged 25-34 constituted 29.2%, while those aged 35-64 represented 14.1%, and those aged 65 and above constituted only 8%.

(Table.1) Socio-demographic features of the study population.



Variable	No. (120)	%
<b>Age (years)</b>		
15-24	67	55.8%
25-34	35	29.2%
35-44	10	8.3%
45-54	3	2.5%
55-64	4	3.3%
65 and above	1	8%
<b>21 Sex</b>		
Male	64	53.3%
Female	56	46.7%
<b>Education</b>		
No education	4	3.3%
Primary school	40	33.3%
Secondary school	50	41.7%
Diploma	11	9.2%
College	12	10.0%
Master	3	2.5%
<b>Marital Status</b>		
Single	85	70.8%
Married	35	29.2%
<b>Residency</b>		
Urban	74	61.7%
Rural	46	38.3%
<b>Is the patient on medical treatment?</b>		
No	7	5.8%
Yes	113	94.2%
<b>Any previous surgical intervention (hip replacement or nerve block)?</b>		
No	69	57.5%
Yes	51	42.5%

Regarding gender distribution, slightly over half of the participants were male (53.3%), while females accounted for 46.7%. In terms of educational status, 41.7% had finished secondary school, 21.7% had attained a higher educational level, and only 3.3% were illiterate. The majority of participants (70.8%) were single, with only 29.2% being married. Additionally, the majority resided in urban areas (61.7%). Regarding disease management, a significant majority (94.2%) were under some form of management, while the remaining 5.8% were not. More than half of the participants (57.5%) had not undergone any surgical intervention, while 42.5% had undergone previous surgical interventions. In terms of disease distribution, the majority (61.67%) had sickle beta thalassemia, while the remainder (38.33%) had sickle cell disease. This is demonstrated in Figure.1.



(Figure.1): Types of Hemoglobinopathies of study population.

Table.2 shows the psycho-social impact of SCD among the study population. Specifically, 81.7% reported feeling happy when interacting with others, while 77.5% expressed concern about experiencing painful episodes during physical activities. Additionally, 74.2% reported feeling irritated or anxious, and 69.2% faced difficulties in their studies or attending school. Eating

limitations were reported by 61.7% of participants, and 57.5% felt restricted in their career choices. Furthermore, 52.5% found it challenging to find a suitable partner for marriage, and 48.3% perceived that SCD negatively impacted their social relationships. About 36.7% did not feel at ease in public settings, and 34.2% lacked a sense of belonging. Fear regarding life expectancy was reported by 55.8% of participants, while 29.2% felt scared and unprotected among friends. Additionally, 23.3% had a history of suicidal thoughts or attempts, and 21.7% felt that people avoided associating with them. Lastly, 18.3% perceived their families as being unconcerned about their well-being.

(Table.2): Psycho-Social impact of SCD among the study population.

Variable	No.	%
<b>SCD adversely impacts my overall social relationships (family, friends)</b>		
No	62	51.7
Yes	58	48.3
<b>My career options are restricted</b>		
No	51	42.5
Yes	69	57.5
<b>I have difficulty in study / get in school</b>		
No	37	30.8
Yes	83	69.2
<b>I struggle to find a compatible partner for marriage/ I face challenges within my marriage</b>		
No	57	47.5
Yes	63	52.5
<b>People avoid being around me</b>		
No	94	78.3
Yes	26	21.7
<b>I feel irritated / anxious</b>		
No	31	25.8
Yes	89	74.2
<b>I feel scared and vulnerable whenever I'm surrounded by friends</b>		
No	85	70.8
Yes	35	29.2
<b>I feel disconnected and lack a sense of belonging</b>		
No	79	65.8
Yes	41	34.2

<b>I feel anxious about my life expectancy</b>		
No	53	44.2
Yes	67	55.8
<b>I feel happy with people</b>		
No	22	18.3
Yes	98	81.7
<b>My family shows no concern for my well-being</b>		
No	98	81.7
Yes	22	18.3
<b>I'm worry about painful episodes during exercises</b>		
No	27	22.5
Yes	93	77.5
<b>Do you have any previous history of suicide or thoughts of suicide?</b>		
No	92	76.7
Yes	28	23.3
<b>I feel unsettled when in public</b>		
No	76	63.3
Yes	44	36.7
<b>I struggle to consume any type of food</b>		
No	46	38.3
Yes	74	61.7


Table.3 presents the average and ranking of respondents' descriptions regarding the psycho-social challenges encountered by sickle cell patients.

(Table.3): Mean and Rank distribution of participants' psychosocial problems facing sickle-cell patient.

<b>Item NO.</b>	<b>Psycho-social problems facing adults with sickle cell disease</b>	<b>Mean</b>	<b>Rank</b>
14	I'm worry about painful episodes during exercises	.78	1st
8	I feel irritated / anxious	.74	2nd
4	I have difficulty in study / get in school	.69	3rd
7	I struggle to consume any type of food	.62	4th
2	My career options are restricted	.58	5th
11	I feel anxious about my life expectancy	.56	6th

3	I struggle to find a compatible partner for marriage/ I face challenges within my marriage	.53	7th
1	SCD adversely impacts my overall social relationships ( family , friends)	.48	8th
6	I feel unsettled when in public	.37	9th
10	I feel disconnected and lack a sense of belonging	.34	10th
9	I feel scared and vulnerable whenever I'm surrounded by friends	.29	11th
15	Do you have any previous history of suicide or thoughts of suicide?	.23	12th
5	People avoid being around me	.22	13th
13	My family shows no concern for my well-being	.18	14th
12	I feel happy with people	.18	15th

Regarding the pain score, the distribution was as follows: 22.50% experienced a little more pain, 21.67% experienced a whole lot of pain, 15.83% experienced a bit of pain, 15.00% experienced even more pain, 13.33% experienced the worst pain, and the lowest percentage, 11.67%, reported no pain. (Figure.2)



(Figure.2): Pain perception score among SCD patient.

In terms of the adverse impact of SCD on social relationships and socio-demographic factors, significant associations were observed with age and pain scores (p-value < 0.05). However, no significant associations were found with other socio-demographic factors such as educational level, sex, and marital status. (Table.4)

(Table.4): The adversely impaction of SCD on social relationships and socio- demographic factors association.

Variables	No		Yes	
	No.	%	No.	%
<b>Age of Respondent(years)</b>				
15-24	39	62.9%	28	55.8%
25 - 34	13	21.0%	22	37.9%
35- 44	6	9.7%	4	6.9%
45 -54	2	3.2%	1	1.7%
55- 64	2	3.2%	2	3.2%
65 and above	0	0.0%	1	1.7%
Total	62	100%	58	100%
Fisher Exact test =5.836		P =0.048		
<b>Education level</b>	<b>No.</b>	<b>%</b>	<b>No.</b>	<b>%</b>
Illiterate	2	3.2%	2	3.2%
Primary education	19	30.6%	21	36.2%
Secondary education	30	48.4%	20	34.5%
Diploma	5	8.1%	6	10.3%
Collage	3	4.8	9	15.5
Master	3	4.8	0	0.0
Total	62	100	58	100
Fisher Exact test =7.735		P =0.065		
<b>Sex</b>	<b>No.</b>	<b>%</b>	<b>No.</b>	<b>%</b>
Male	31	50.0	33	53.3
Female	31	50.0	25	46.7
X <sup>2</sup> =1.503		DF = 1	P = 0.110	
<b>Marital Status</b>	<b>No.</b>	<b>%</b>	<b>No.</b>	<b>%</b>

Single	45	72.6	40	69.0
Married	17	27.4	18	29.2
$X^2 = 0.190$		DF = 1	P = 0.145	
<b>Pain Score</b>	<b>No.</b>	<b>%</b>	<b>No.</b>	<b>%</b>
No Hurt	8	12.9	6	11.7
Hurts little bit	10	16.1	9	15.8
Hurts little more	11	17.7	16	27.6
Hurts even more	14	22.6	4	6.9
Hurts whole lot	13	21.0	13	22.4
Hurts worst	6	9.7	10	13.3
$X^2 = 7.695$		DF = 5	P = 0.04	

## Discussion

Apart from the physical health issues associated with sickle cell disease, patients also face complex psychosocial obstacles, including pain crises, sleep disorders, interpersonal relationships and social stigma. These challenges also impact their families and social networks, and can significantly affect the overall effectiveness of treatment [10]. This cross-sectional study involving 120 SCD patient, revealed <sup>25</sup> that the majority of cases fell within the age group of 15 to 24 years old. More than half of the participants were male (53.3%), while the remaining (46.7%) were female. Additionally, the majority of participants (70.8%) were single, with only 29.2% being married. Similar <sup>1</sup> to a previous study conducted in Nigeria by Theresa et al. [16], the majority of respondents (61.7%) hailed from urban areas. This also align with the Sickle Cell Data Collection Program (SCDC) in Tennessee by Allison et al. [17]. <sup>1</sup> In terms of educational level, a similar distribution was observed by Harris et al. [18], with the vast majority of patients had a secondary school degree or a higher educational degree.

Regarding management, the majority of patients (94.2%) were undergoing treatment, while the remaining 5.8% were not. These results are in line with those obtained <sup>5</sup> from the international

Sickle Cell World Assessment Survey (SWAY) by Osunkwo et al. [19]. Furthermore, more than half of the participants (57.5%) had not undergone any procedures like hip replacements or nerve blocks, while 42.5% had undergone previous surgical interventions. A similar trend was seen in a cross-sectional study by Alshurafa et al. [20].

Moreover, 57.5% reported limitations in their choice of career, which has severe psychosocial implications, similar to findings from a previous study by Thomas et al. [21] Additionally, a notably high proportion of patients (74.2%) felt irritated or anxious, consistent with findings suggesting increased depression and anxiety among individuals with SCA by Molock et al. [22]. Additionally, concerns about painful episodes during exercise were reported by 77.5% of participants by Gouraud et al. [23] concluded that, in addition to the effect of anemia in SCD on diminishing oxygen delivery to the skeletal muscle, patients also exhibit significant skeletal muscle dysfunction, resulting in reduction in functional capacity and overall quality of life. Furthermore, 48.3% perceived that SCA adversely affected their overall social relationships. This is consistent with the findings of Adzika et al. [24].

Difficulties in studying or attending school were reported by 69.2% of participants. A previous cross-sectional survey of SCA patients by Osunkwo et al. showed significant impact of SCD on achievement in school and career progress a finding which is consistent with the present study(19) Additionally, 52.5% found it challenging to find a suitable partner in marriage. Oladiran et al. [25] studied the influence of SCD on the choice of romantic partners. They reported concerns about passing on the disease, caregiving responsibilities, lifestyle limitations, fear of complications, and stigma surrounding SCD. Moreover, 23.3% reported a history of suicide or suicidal thoughts, consistent with a previous study by Oliveira et al. who reported a percentage of 20%(26). This could be attributed to the strong religious beliefs among the population, as religion has been linked to lower rates of suicidal thoughts in some studies [27].



Regarding the association between the adverse impacts of SCA on social relationships and socio-demographic factors, there was a significant association with age and pain scores, with a p-value less than 0.05, similar to findings from previous studies by Edwards et al. [28] and McClish et al. [29]. However, no significant associations were found with other socio-demographic factors such as educational level, sex, and marital status. This aligns with findings from studies by Harris et al. [18] and Adzika et al. [24].

### **Limitations of the study**

While the study provided valuable insights into the targeted population, several limitations need consideration. These include the potential response bias in self-reported data, the cross-sectional nature limiting causal inferences, subjective pain assessment methods, and limited sample size. Despite these limitations, the study contributes to understanding the psychosocial challenges faced by SCD patients, highlighting the need for further research addressing these complexities in diverse settings and populations.

### **Conclusion**

This study highlights the significant psychosocial challenges faced by SCD patients, affecting various aspects of their lives including education, employment, relationships, and mental health. While certain socio-demographic factors showed associations with pain scores, further research is needed to better understand the complex interplay between these factors and the impact of SCD.

**Funding and sponsorship:** Nil.

**Conflicts of interest:** No conflicts of interest were present in this study.

## REFERENCES

1. Payne AB, Mehal JM, Chapman C, et al. Trends in sickle cell disease–related mortality in the United States, 1979 to 2017. *Annals of Emergency Medicine*. 2020;76(3):S28-S36.
2. Kato G, Piel F, Reid C, et al. Sickle cell disease. *Nat Rev Dis Primers*. 2018;4:18010.
3. Knisely MR, Pugh N, Kroner B, et al. Patient-reported outcomes in sickle cell disease and association with clinical and psychosocial factors: Report from the sickle cell disease implementation consortium. *American Journal of Hematology*. 2020;95(9):1066-74.
4. Brandow AM, Carroll CP, Creary S, et al. American Society of Hematology 2020 guidelines for sickle cell disease: management of acute and chronic pain. *Blood Advances*. 2020;4(12):2656-701.
5. Sil S, Cohen LL, Dampier C. Psychosocial and functional outcomes in youth with chronic sickle cell pain. *The Clinical Journal of Pain*. 2016;32(6):527-33.
6. Sagi V, Mittal A, Tran H, et al. Pain in sickle cell disease: current and potential translational therapies. *Translational Research*. 2021;234:141-58.
7. Osunkwo I, O'Connor HF, Saah E. Optimizing the management of chronic pain in sickle cell disease. *Hematology 2014, the American Society of Hematology Education Program Book*. 2020;2020(1):562-9.
8. Takaoka K, Cyril AC, Jinesh S, et al. Mechanisms of pain in sickle cell disease. *British Journal of Pain*. 2021;15(2):213-20.
9. Brandow A, Liem R. Advances in the diagnosis and treatment of sickle cell disease. *Journal of Hematology & Oncology*. 2022;15(1):20.
10. Essien EA, Winter-Eteng BF, Onukogu CU, et al. Psychosocial challenges of persons with sickle cell anemia: A narrative review. *Medicine*. 2023;102(47):e36147.
11. Adam SS, Flahiff CM, Kamble S, et al. Depression, quality of life, and medical resource utilization in sickle cell disease. *Blood Advances*. 2017;1(23):1983-92.
12. Bulgin D, Tanabe P, Jenerette C. Stigma of sickle cell disease: a systematic review. *Issues in Mental Health Nursing*. 2018;39(8):675-86.
13. Valrie CR, Alston K, Fuh B, et al. Sleep moderating the relationship between pain and health care use in youth with sickle cell disease. *The Clinical Journal of Pain*. 2020;36(2):117-23.

14. Gillespie ML, Spring MR, Cohen RT, et al. The interplay of sleep disordered breathing, nocturnal hypoxemia, and endothelial dysfunction in sickle cell disease. *Progress in Pediatric Cardiology*. 2023;68:101602.
15. Al-Marzouki AF, Alrefaie NI, Aljohani NA, et al. The prevalence of depression and anxiety among sickle cell disease patients in King Abdulaziz University Hospital. *Cureus*. 2021;13(9).
16. Nwagha T, Omotowo BI. Determinants of psychosocial health-related quality of life of adults with sickle cell disease in a Nigerian setting. *Nigerian Medical Journal: Journal of the Nigeria Medical Association*. 2020;61(3):114.
17. Plaxco AP, Hankins JS, Davis R, et al. Descriptive epidemiology of sickle cell disease in Tennessee: population-based estimates from 2008 to 2019. *Frontiers in Hematology*. 2023;2:1277548.
18. Harris KM, Preiss L, Varughese T, et al. Examining mental health, education, employment, and pain in sickle cell disease. *JAMA Network Open*. 2023;6(5):e2314070-e.
19. Osunkwo I, Andemariam B, Minniti CP, et al. Impact of sickle cell disease on patients' daily lives, symptoms reported, and disease management strategies: Results from the international Sickle Cell World Assessment Survey (SWAY). *American Journal of Hematology*. 2021;96(4):404-17.
20. Alshurafa A, Soliman AT, De Sanctis V, et al. Clinical and epidemiological features and therapeutic options of avascular necrosis in patients with sickle cell disease (SCD): A cross-sectional study. *Acta Bio Medica: Atenei Parmensis*. 2023;94(5).
21. Thomas VJ, Taylor L. The psychosocial experience of people with sickle cell disease and its impact on quality of life: Qualitative findings from focus groups. *British Journal of Health Psychology*. 2002;7(3):345-63.
22. Molock SD, Belgrave FZ. Depression and anxiety in patients with sickle cell disease: conceptual and methodological considerations. *Journal of Health & Social Policy*. 1994;5(3-4):39-53.
23. Gouraud E, Connes P, Gauthier-Vasserot A, et al. Is skeletal muscle dysfunction a limiting factor of exercise functional capacity in patients with sickle cell disease? *Journal of Clinical Medicine*. 2021;10(11):2250.

24. Adzika VA, Glozah FN, Ayim-Aboagye D, et al. Socio-demographic characteristics and psychosocial consequences of sickle cell disease: the case of patients in a public hospital in Ghana. *Journal of Health, Population and Nutrition*. 2017;36:1-10.
25. Oladiran O-O. Exploring How Sickle Cell Disease Impacts the Selection of Romantic Partner and Reproductive Decision Making of Adults with Sickle Cell Disease Living in The United Kingdom: University of Sheffield; 2021.
26. de Oliveira LAB, da Silva Lopes T, de Sousa JN, et al. Religiosity, Anxiety, Depression, and Suicidal Ideation in Brazilian Patients with Sickle Cell Disease. *REFLEXUS-Revista Semestral de Teologia e Ciências das Religiões*. 2023;17(2):277-90.
27. Hajjiyousouf II, Bulut S. Mental health, religion and suicide. *Open Journal of Medical Psychology*. 2021;11(1):12-27.
28. Edwards CL, Killough A, Wood M, et al. Emotional reactions to pain predict psychological distress in adult patients with Sickle Cell Disease (SCD). *The International Journal of Psychiatry in Medicine*. 2014;47(1):1-16.
29. McClish DK, Smith WR, Levenson JL, et al. Comorbidity, pain, utilization, and psychosocial outcomes in older versus younger sickle cell adults: the PiSCES project. *BioMed Research International*. 2017;2017.