

Study to estimate the NIHS score and volume of stroke in predicting 30 day mortality in acute ischemic stroke patients

By Dhanush Balaji S

Type of the Manuscript: Research article

Study to estimate the NIHSS score and volume of stroke in predicting 30 day mortality in acute ischemic stroke patients

Dhanush Balaji S.¹, Abinaya Srinivasa Rangan¹, S. Prasanna Karthik¹, Karpaka Vinayakam Gopalakrishnan², Selva Balaji¹ Manoj Kumar S¹

¹Department Of General Medicine, Saveetha Medical College and Hospital , Tamil Nadu

² Saveetha Medical College and Hospital, Saveetha Institute of Medical and Technical Sciences (SIMATS), Saveetha University, Chennai, India

Corresponding author:

Dhanush Balaji.S

E-mail id: dhanushbalajis@gmail.com

ABSTRACT

Background: Stroke is defined as a sudden onset of a neurological deficit attributable to a focal vascular cause. It is one of the leading causes of mortality and morbidity all over the world. Present study was aimed to estimate the NIHSS score and volume of stroke in predicting 30 day mortality in acute ischemic stroke patients.

Material and methods: Present study was prospective cross sectional study, conducted in patients with age more than 18 years, either gender, symptoms of stroke less than 24 hours, diagnosis of acute ischemic stroke based on history, physical examination, CT scan and/or diffusion weighted MRI scan performed during the first 24 hours.

Results: In present study, mean age was 57.4 ±13.36 years. The volume of stroke on 1st day was 40.75 ±38.51 and on 30th day it reduced to 31.94 ±33.20 which was statistically significant. NIHSS (National Institutes of Health Stroke Scale) score on day 1 was 9.70 ±5.22 and on day 30 the score was 7.77±5.25, difference was statistically significant (p=0.000). At end of 30 days, 10 % mortality was observed in present study. NIHSS score & volume of stroke were significantly associated with 30 days outcome. **Conclusion:** Volume of stroke by MRI brain and NIHSS scoring are well correlated in predicting 30 day mortality in acute ischemic stroke patients.

Keywords: stroke, NIHSS, volume of stroke, MRI brain, acute ischemic stroke

INTRODUCTION

The abrupt onset of a neurological loss resulting from a focal vascular source is referred to as a stroke [1]. Across the globe, it is one of the main causes of both mortality and morbidity. There are two types of stroke: ischemic and hemorrhagic. 15% of strokes are hemorrhagic, while 85% of all strokes are ischemic [2]. There are several risk factors for stroke that are discussed, including high blood pressure, a history of stroke or transient ischemic attack, high cholesterol, heart disease, diabetes, and sickle cell disease. Stroke was also linked to lifestyle-related factors such as smoking, drinking too much alcohol, obesity, poor diet, and lack of physical activity [3]. The Glasgow Coma Scale (GCS), Canadian Neurological Scale (CNS), and National Institute of Health Stroke Scale (NIHSS) are used to evaluate the prognosis of acute ischemic stroke. Because of its rigorous design and simplicity, NIHSS is widely acknowledged [4]. It has been shown to be a highly accurate predictor of patient outcomes, evaluating patients based on eleven distinct characteristics. One important factor influencing prognosis in acute stroke patients is the quantity of brain tissue that has had reversible damage at the time of presentation. A worse prognosis and spontaneous hemorrhagic change are often linked to larger infarcts in ischemic stroke [5]. The purpose of the current study was to estimate the NIHSS score and stroke volume in order to predict the 30-day death rate for patients with acute ischemic stroke.

MATERIAL AND METHODS

This prospective cross-sectional study was carried out at Saveetha Medical College and Hospital's general medicine department in Chennai, India. The two-year study period ran from January 2021 until December 2022. It was approved by the institutional ethical committee to conduct the study. Patients who meet the following criteria will be included in the study: they must be over the age of 18, male or female, have symptoms of a stroke within 24 hours, have been diagnosed with acute ischemic stroke based on history, physical examination, CT scan or diffusion weighted MRI scan completed within the first 24 hours, and be willing to participate in the study.

Exclusion criteria: Intracerebral mass, subdural hematoma, and cerebral hemorrhage diagnosed by CT scan. Individuals that exhibit hemodynamic stability and are unsuitable for MRI testing for a variety of reasons. Individuals with a prior history of stroke were known to have hematological problems that impact red blood cells and platelets.

Individuals unwilling to provide written informed consent.

Patients were informed about the study in their native tongue. The stroke patient's family gave their signed, informed consent. Patient information was documented, including demographics, medical history, co-morbidities, drug and alcohol use, alcohol intake, smoking and nicotine use, trauma, and previous history of TIA or stroke. All patients underwent a thorough clinical examination upon admission to assess neurological impairment, vital statistics, and Glasgow Coma Scale scores.

All patients had routine tests, including full hemogram, ESR, BT, PT, aPTT, platelet count, routine and microscopic urine examination, serum creatinine, eGFR, serum electrolytes, LFT, lipid profile, CT scan/MRI head, ECG, and chest X-ray. Whenever necessary, additional tests were carried out, including connective tissue workups and echocardiograms.

Every patient got the usual care. The assessment of stroke patients' outcome was based on their mortality rate 30 days following the stroke occurrence. Follow-up continued for three months. Day 1 and Day 30 NIHSS scores and the volume of stroke (based on MRI) were computed.

SPSS 23.0 was used for data analysis after Microsoft Excel was used for data collection and compilation. For the continuous variables, frequency, percentage, means, and standard deviations (SD) were computed; for the categorical variables, ratios and proportions were computed. The chi-square test or the Fisher exact test, if appropriate, were used to evaluate the differences in proportions between the qualitative variables. A P-value of less than 0.5 was deemed statistically noteworthy.

RESULTS

In present study, mean age was 57.4 ± 13.36 with the minimum age of 24 years and maximum of 83 years. Majority patients were between 41-60 years of age (44 %), followed by above 60 years age (42 %). 55% of our subjects were males in our study. The most common co-morbidities were diabetes mellitus (46 %), followed by hypertension (27 %), sustained hypertension (21 %) & chronic kidney diseases (6 %). [TABLE 1]. The volume of stroke on 1st day was 40.75 ± 38.51 and on 30th day it reduced to 31.94 ± 33.20 which was statistically significant. NIHSS (National Institutes of Health Stroke Scale) score on day 1 was 9.70 ± 5.22 and on day 30 the score was 7.77 ± 5.25 , difference was statistically significant ($p=0.000$). [TABLE 2]

Severity of stroke based on NIHSS score on 1st day of acute ischemic stroke patients was 3.50 ± 1.75 , moderate category score was 29.46 ± 8.417 , moderate to severe stroke was 81.56 ± 14.48 and severe stroke category was 161.20 ± 38.51 . Intergroup comparison by post Hocktuckey test showed

statistical results in between all groups. [TABLE 3]

NIHSS score on 30th day of acute ischemic stroke patients, in minor stroke category was 3.47 ± 2.07 , 31.79 ± 11.18 was score of moderate category, score in moderate to severe category was 80.43 ± 9.93 and score in severe category was 160.40 ± 6.150 . The intergroup comparison was highly statistically significant ($p=0.000$). [TABLE 3]

The volume of stroke in mild, moderate, moderate to severe and severe stroke category on day 1 was 3.50 ± 1.75 , 29.46 ± 8.417 , 81.56 ± 14.48 and 161.20 ± 38.51 respectively. The volume of stroke in mild, moderate, moderate to severe and severe stroke category on day 30 was 3.47 ± 2.07 , 31.79 ± 11.18 , 80.43 ± 9.93 and 160.40 ± 6.150 respectively. Except in moderate category all other category had decrease in the volume of stroke on day 30 compared to day 1. Though the results are not significant in all categories except moderate, the volume of stroke was reduced on day 30.

NIHSS score decreased on day 30 in moderate and moderate to severe category and the score is increased slightly in mild and severe category. This shows that the overall severity of our subjects were greatly reduced and it was statistically significant ($p=0.000$). [TABLE 4]

At end of 30 days, 10 % mortality was observed in present study. NIHSS score & volume of stroke were significantly associated with 30 days outcome. [TABLE 5]

DISCUSSION

Ischemic stroke is brought on by a disruption in the blood supply to the brain. Many clinical and medical factors, including age, comorbid conditions, sex, the size and severity of the infarct, the aetiology and location of the stroke, the amount of time that passes between the stroke's onset and hospitalization, and the type of medical care provided, such as stroke unit, stroke team, and stroke pathway, all affect the prognosis for functional recovery following a stroke [6].

In our study, we have observed that the mean infarct volume of stroke on 1st day was 40.75 ± 38.51 and on 30th day it reduced to 31.94 ± 33.20 which was statistically highly significant ($p=0.000$). Infarct volume is one of the important predictors of mortality in acute ischemic stroke. Reduction in infarct volume has

been encouraged as a substitute²⁵ outcome measure for human stroke clinical trials. Past studies have shown correlations between subacute CT lesion size and clinical scales such as NIHSS, Rankin Disability Scale, Oxford Disability Scale and aphasia severity scale [7]. Within the mild, moderate, moderate to severe, and severe stroke categories in our study, the volume of strokes on day 1 was 3.50 ± 1.75 , 29.46 ± 8.417 , 81.56 ± 14.48 , and 161.20 ± 38.51 , respectively. On day 30, the number of stroke cases in the mild, moderate, moderate to severe, and severe stroke categories was 3.47 ± 2.07 , 31.79 ± 11.18 , 80.43 ± 9.93 , and 160.40 ± 6.150 , in that order. On day 30, there was a significant decrease in the volume of stroke in all categories except moderate. There was a substantial statistical difference between the groups ($p=0.000$).

We have seen a strong correlation between individuals' NIHSS scores and the amount of stroke. Given the relationship between CT infarct volume and accepted clinical outcome indicators, CT volume may be used as a prognostic indicator to estimate 30-day mortality in cases of acute ischemic stroke. Although the NIH Stroke Scale (NIHSS) was initially intended for prospective scoring, it is now utilized to determine the severity of a stroke retrospectively [8,9]. Baseline NIHSS scores that are highly correlated with lesion volume show a strong likelihood of mortality for scores greater than 16, and a strong likelihood of a good recovery for scores less than 6 [10].

The NIHSS score was 9.70 ± 5.22 on day 1 and 7.77 ± 5.25 on day 30. At $p = 0.000$, this is statistically significant. Patients with acute ischemic stroke can be accurately predicted by having elevated NIHSS scores. According to our analysis, three of them fell into the category of severe stroke after 30 days. Of the 3,2 that passed away, 2 had complications from an acute ischemic stroke, meaning that 66.7 percent of patients in the severe group, as defined by the NIHSS, were at danger of dying. In our investigation, we found a strong correlation between the volume of stroke and NIHSS. Significant correlation was found at the 0.01 (2-tailed) level. The development of stroke and its effects may be prevented by early diagnosis, treatment, including lifestyle change, and prevention of diabetes. This is a significant challenge for medical professionals dealing with the epidemic of both diabetes and stroke. Future research should evaluate the precise reasons for the poor result and whether a more aggressive treatment strategy will enhance the prognosis of these patients.

CONCLUSION

We found a significant connection between the volume of stroke and NIHSS at the 0.01 level. As a result, in acute ischemic stroke patients, the volume of stroke by MRI brain and the NIHSS scoring have a strong correlation when it comes to predicting 30-day mortality.

4
Author Contributions: Conceptualization, D.S. and P.K.; methodology, D.S.; software, D.S. and K.G.; validation, D.S., A.S. and M.K.; formal analysis, D.S.; investigation, D.S.; resources, D.S.; data curation, D.S.; writing—original draft preparation, D.S., P.K.; writing—review and editing, D.S., K.G.; abstract - S.A.; visualization, D.S., A.S.; supervision, P.K.; project administration, D.S. All authors have read and agreed to the published version of the manuscript.

Conflicts of Interest: The authors declare no conflicts of interest

Funding: This research received no external funding



TABLES

Table 1- General characteristics and prevalence of various co-morbidities

Characteristics	No. of patients	Percentage
Age groups (in years)		
20-40	14	14
41-60	44	44
>60	42	42
Mean age (mean \pm SD)	57.4 \pm 13.36	
Gender		
Male	55	55
Female	45	45
Co-morbidities		
Type 2 Diabetes Mellitus	46	46
Hypertension	27	27
Sustained Hypertension	21	21
Chronic Kidney Diseases	06	06

Table 2: Comparison of 1st day and 30th day parameters

Parameter	on 1 st day	on 30 th day	Mean difference	t-value	p value
Volume of stroke	40.75 \pm 38.51	31.9 \pm 33.20	8.818	5.200	0.000***
NIHSS Score	9.70 \pm 5.22	7.77 \pm 5.25	1.93	8.36	0.000***

Table 3: Comparison of Volume of Stroke in severity of stroke on 1st day and on 30th day

Severity of Stroke based on NIHSS Score (N)	Volume of Stroke on 1 st day	F-Value	p-value	Intergroup Comparison by post Hoc Tukey test
Minor Stroke (20)	3.50 ± 1.75	570.42	0.000***	1-2-0.000***
Moderate Stroke (56)	29.46 ± 8.417			1-3-0.000***
Moderate–Severe Stroke (19)	81.56 ± 14.48			1-4-0.000***
Severe Stroke (05)	161.20 ± 38.51			2-3-0.000***
				2-4-0.000***
				3-4-0.000***
Severity of Stroke based on NIHSS Score (N)	Volume of Stroke on 30 th day	F-Value	p-value	Intergroup Comparison by post Hoc Tukey test
Minor Stroke (20)	3.47 ± 2.07	414.126	0.000***	1-2-0.000***
Moderate Stroke (56)	31.79 ± 11.18			1-3-0.000***
Moderate–Severe Stroke (19)	80.43 ± 9.93			1-4-0.000***
Severe Stroke (05)	160.40 ± 6.150			2-3-0.000***
				2-4-0.000***
				3-4-0.000***

Table 4: Comparison of Grade of Stroke severity between 1st day and 30th day

Stroke Severity on 1 st day	Stroke Severity on 30 th day				Total
	Minor Stroke N(%)	Moderate Stroke N(%)	Moderate–Severe Stroke N(%)	Severe Stroke N(%)	
Minor Stroke	20 (20)	0	0	0	20 (20)
Moderate Stroke	12 (12)	44 (44)	0	0	56 (56)
Moderate–Severe Stroke	0	10 (10)	09 (09)	0	19 (19)
Severe Stroke	0	0	02 (02)	03 (03)	05 (05)
Total	32(32)	54 (54)	11 (11)	03 (03)	100 (100)
Fisher Exact Test:-X2value=100.523,p-Value=0.000***					

Table 5: Comparison of parameters among survivor and death patients

Parameter	Survivor patients N=90		Death Patients N=10	
	Parameter at 1 st Day Mean±SD	Parameter at 30 th Day Mean±SD	Parameter at 1 st Day Mean±SD	Parameter at 30 th Day Mean±SD
NIHSS score	8.28±3.90	6.44±3.92	19.80±1.75	19.20±2.04
	tvalue:8.727;pvalue: 0.000***		tvalue:2.714;pvalue:0.024**	
Volume of Stroke	32.19±25.27	25.28±21.95	132.16±34.61	100.13±50.67
	tvalue:4.724;pvalue: 0.000***		tvalue:2.855;pvalue:0.02**	

*p<0.05- **p<0.001- statistically significant, ns-not significant